



Bioenergetics

Cumulative Contents

Rapid report

Direct evidence for excitonically coupled chlorophylls *a* and *b* in LHC II of higher plants by nonlinear polarization spectroscopy in the frequency domain

M. Krikunova, B. Voigt and H. Lokstein (Germany, Russia)

1

Regular papers

Effect of anoxia/reperfusion on the reversible active/de-active transition of NADH-ubiquinone oxidoreductase (complex I) in rat heart

E. Maklashina, Y. Sher, H.-Z. Zhou, M.O. Gray, J.S. Karliner and G. Cecchini (USA)

6

Photoaccumulation of the PsbB phytylsemiquinone in Photosystem I of *Chlamydomonas reinhardtii*

S.E.J. Rigby, I.P. Muhiuddin, M.C.W. Evans, S. Purton and P. Heathcote (UK)

13

Photosystem II reaction center with altered pigment composition: reconstitution of a complex containing five chlorophyll *a* per two pheophytin *a* with modified chlorophylls

A. Zehetner, H. Scheer, P. Siffel and F. Vacha (Germany, Czech Republic)

21

The Lhca antenna complexes of higher plants photosystem I

R. Croce, T. Morosinotto, S. Castelletti, J. Breton and R. Bassi (Italy, France)

29

Mitochondrial respiratory rates and activities of respiratory chain complexes correlate linearly with heteroplasmy of deleted mtDNA without threshold and independently of deletion size

F.N. Gellerich, M. Deschauer, Y. Chen, T. Müller, S. Neudecker and S. Zierz (Germany)

41

Proteolytic activity against the light-harvesting complex and the D1/D2 core proteins of Photosystem II in close association to the light-harvesting complex II trimer

J.H. Georgakopoulos, A. Sokolenko, M. Arkas, G. Sofou, R.G. Herrmann and J.H. Argyroudi-Akoyunoglou (Greece, Germany)

53

Full recovery of the NADH:ubiquinone activity of complex I (NADH:ubiquinone oxidoreductase) from *Yarrowia lipolytica* by the addition of phospholipids

S. Dröse, K. Zwicker and U. Brandt (Germany)

65

Characterization of the mitochondrial respiratory pathways in *Candida albicans*

E.J. Helmerhorst, M.P. Murphy, R.F. Troxler and F.G. Oppenheim (USA, United Kingdom)

73

Orientation selection in photosynthetic PS I multilayers: structural investigation of the charge separated state $P_{700}^{+}A_1^{-}$ by high-field/high-frequency time-resolved EPR at 3.4 T/95 GHz

M. Fuhs, A. Schnegg, T. Prisner, I. Köhne, J. Hanley, A.W. Rutherford and K. Möbius (Germany, France)

81

Rapid reports

Raman spectroscopy of chlorophyll *d* from *Acaryochloris marina*

Z.-L. Cai, H. Zeng, M. Chen and A.W.D. Larkum (Australia)

89

Introduction of the $^{305}\text{Arg} \rightarrow ^{305}\text{Ser}$ mutation in the large extrinsic loop E of the CP43 protein of *Synechocystis* sp. PCC 6803 leads to the loss of cytochrome c_{550} binding to Photosystem II

T.M. Bricker, A. Young, L.K. Frankel and C. Putnam-Evans (USA)

92

Regular papers

Opening of potassium channels modulates mitochondrial function in rat skeletal muscle

G. Debska, A. Kicinska, J. Skalska, A. Szewczyk, R. May, C.E. Elger and W.S. Kunz (Poland, Germany)

97

Hybrid ubiquinone: novel inhibitor of mitochondrial complex I

H. Yabunaka, A. Kenmochi, Y. Nakatogawa, K. Sakamoto and H. Miyoshi (Japan)

106

Kinetics of the spectral changes during reduction of the Na ⁺ -motive NADH:quinone oxidoreductase from <i>Vibrio harveyi</i> <i>A.V. Bogachev, Y.V. Bertsova, E.K. Ruuge, M. Wikström and M.I. Verkhovsky (Russia, Finland)</i>	113	The H ⁺ -ATPase from chloroplasts: effect of different reconstitution procedures on ATP synthesis activity and on phosphate dependence of ATP synthesis <i>I. Grotjohann and P. Gräber (Germany)</i>	208
Transmembrane topology of the NuoL, M and N subunits of NADH:quinone oxidoreductase and their homologues among membrane-bound hydrogenases and bona fide antiporters <i>C. Mathiesen and C. Høgerhäll (Sweden)</i>	121	The NADP-reducing hydrogenase from <i>Desulfovibrio fructosovorans</i> : functional interaction between the C-terminal region of HndA and the N-terminal region of HndD subunits <i>Z. Dermoun, G. De Luca, M. Asso, P. Bertrand, F. Guerlesquin and B. Guigliarelli (France)</i>	217
Dimerization of F ₀ F ₁ ATP synthase from bovine heart is independent from the binding of the inhibitor protein IF ₁ <i>L. Tomasetig, F. Di Pancrazio, D.A. Harris, I. Mavelli and G. Lippe (Italy, UK)</i>	133	Nonoxidizable ubiquinol derivatives that are suitable for the study of the ubiquinol oxidation site in the cytochrome bc ₁ complex <i>L. Zhang, Z. Li, B. Quinn, L. Yu and C.-A. Yu (USA)</i>	226
Expression of <i>chaA</i> , a sodium ion extrusion system of <i>Escherichia coli</i> , is regulated by osmolarity and pH <i>T. Shijuku, T. Yamashino, H. Ohashi, H. Saito, T. Kakegawa, M. Ohta and H. Kobayashi (Japan)</i>	142	Mitochondrial damage by nitric oxide is potentiated by dopamine in PC12 cells <i>F. Antunes, D. Han, D. Rettori and E. Cadenas (USA, Portugal)</i>	233
Squalane is in the midplane of the lipid bilayer: implications for its function as a proton permeability barrier <i>T. Hauf, S. Dante, N.A. Dencher and T.H. Haines (Germany, USA)</i>	149	Resolution of the Photosystem I and Photosystem II contributions to chlorophyll fluorescence of intact leaves at room temperature <i>F. Franck, P. Juneau and R. Popovic (Belgium, Canada)</i>	239
Bax and heart mitochondria: uncoupling and inhibition of respiration without permeability transition <i>F. Appaix, K. Guerrero, D. Rampal, M. Izikki, T. Kaambre, P. Sikk, D. Brdiczka, C. Riva-Lavieille, J. Olivares, M. Longuet, B. Antonsson and V.A. Saks (France, Estonia, Germany, Switzerland)</i>	155	<i>Rhodopseudomonas acidophila</i> strain 10050 contains photosynthetic LH2 antenna complexes that are not enriched with phosphatidylglycerol, and the phospholipids have a fatty acyl composition that is unusual for purple non-sulfur bacteria <i>N.J. Russell, J.K. Coleman, T.D. Howard, E. Johnston and R.J. Cogdell (UK)</i>	247
Refined structure of c-phycoerythrin from the cyanobacterium <i>Synechococcus vulcanus</i> at 1.6 Å: insights into the role of solvent molecules in thermal stability and co-factor structure <i>N. Adir, R. Vainer and N. Lerner (Israel)</i>	168	Stabilization of iron–sulfur cluster F _x by intra-subunit interactions unraveled by suppressor and second site-directed mutations in PsaB of Photosystem I <i>M.-T. Zeng, X.-M. Gong, M.C.W. Evans, N. Nelson and C. Carmeli (Israel, USA, UK)</i>	254
Spectroscopic and oxidation–reduction properties of <i>Rhodobacter capsulatus</i> cytochrome c ₁ and its M183K and M183H variants <i>J. Li, E. Darrouzet, I.K. Dhawan, M.K. Johnson, A. Osyczka, F. Daldal and D.B. Knaff (USA)</i>	175	Spectroscopic properties of PSI–IsiA supercomplexes from the cyanobacterium <i>Synechococcus</i> PCC 7942 <i>E.G. Andrizhiyevskaya, T.M.E. Schwabe, M. Germano, S. D'Haene, J. Kruip, R. van Grondelle and J.P. Dekker (The Netherlands, Germany)</i>	265
Interaction of genistein with the mitochondrial electron transport chain results in opening of the membrane transition pore <i>M. Salvi, A.M. Brunati, G. Clari and A. Toninello (Italy)</i>	187	Direct electron transfer between hemoglobin and a glassy carbon electrode facilitated by lipid-protected gold nanoparticles <i>X. Han, W. Cheng, Z. Zhang, S. Dong and E. Wang (China)</i>	273
Relevance of the diastereotopic ligation of magnesium atoms of chlorophylls in Photosystem I <i>T.S. Balaban, P. Fromme, A.R. Holzwarth, N. Krauß and V.I. Prokhorenko (Germany, USA)</i>	197	<i>Author Index</i>	278
		<i>Cumulative Contents</i>	281